



HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843

Tel: 714-891-2022; Fax: 714-891-2123

www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Status: FINAL REPORT
Notes: hx of food intolerance, scratching, soft stool
Diet: Raw Diet

Accession No. Test10000	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 08/23/18
Species Canine	Breed Golden Retriever	Sex FS	Pet Age 5 Yrs 45 lbs	Reported 08/23/18

Test Requested	Result	Case Specific	General Range	Units
Beef Salivary IgA	12.850 Intermediate reaction, Avoid		< 10	U/mL
Beef Salivary IgM	8.600 Negative Reaction		< 10	U/mL
Chicken Salivary IgA	12.518 Intermediate reaction, Avoid		< 10	U/mL
Chicken Salivary IgM	10.344 Weak Reaction		< 10	U/mL
Corn Salivary IgA	11.316 Weak Reaction		< 10	U/mL
Corn Salivary IgM	11.607 Borderline Reaction; Avoid		< 10	U/mL
Duck Salivary IgA	10.401 Weak Reaction		< 10	U/mL
Duck Salivary IgM	10.695 Weak Reaction		< 10	U/mL
Lamb Salivary IgA	9.495 Negative Reaction		< 10	U/mL
Lamb Salivary IgM	10.184 Weak Reaction		< 10	U/mL
Milk Salivary IgA	10.809 Weak Reaction		< 10	U/mL
Milk Salivary IgM	10.926 Weak Reaction		< 10	U/mL
Pork Salivary IgA	10.475 Weak Reaction		< 10	U/mL
Pork Salivary IgM	10.350 Weak Reaction		< 10	U/mL
Soy Salivary IgA	10.633 Weak Reaction		< 10	U/mL
Soy Salivary IgM	9.727 Negative Reaction		< 10	U/mL
Turkey Salivary IgA	10.238 Weak Reaction		< 10	U/mL
Turkey Salivary IgM	11.159 Weak Reaction		< 10	U/mL
Venison Salivary IgA	12.172 Intermediate reaction, Avoid		< 10	U/mL
Venison Salivary IgM	10.901 Weak Reaction		< 10	U/mL
Wheat Salivary IgA	11.236 Weak Reaction		< 10	U/mL
Wheat Salivary IgM	10.799 Weak Reaction		< 10	U/mL
White Fish Salivary IgA	11.980 Borderline Reaction; Avoid		< 10	U/mL



ENRICHING PET LIFE THROUGH DIAGNOSTIC TESTING
world-wide personal mail-in diagnostic lab for vets and pets



HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843

Tel: 714-891-2022; Fax: 714-891-2123

www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Accession No. Test10000	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 08/23/18
-----------------------------------	-------------------------	-----------------------------------	---------------------------	----------------------

Test Requested	Result	Case Specific	General Range	Units
White Fish Salivary IgM	11.140 Weak Reaction		< 10	U/mL
Barley Salivary IgA	10.200 Weak Reaction		< 10	U/mL
Barley Salivary IgM	9.800 Negative Reaction		< 10	U/mL
Egg Salivary IgA	8.680 Negative Reaction		< 10	U/mL
Egg Salivary IgM	8.470 Negative Reaction		< 10	U/mL
Lentil Salivary IgA	10.050 Weak Reaction		< 10	U/mL
Lentil Salivary IgM	12.257 Intermediate reaction, Avoid		< 10	U/mL
Millet Salivary IgA	9.266 Negative Reaction		< 10	U/mL
Millet Salivary IgM	10.275 Weak Reaction		< 10	U/mL
Oatmeal Salivary IgA	9.900 Negative Reaction		< 10	U/mL
Oatmeal Salivary IgM	12.350 Intermediate reaction, Avoid		< 10	U/mL
Peanut Salivary IgA	10.306 Weak Reaction		< 10	U/mL
Peanut Salivary IgM	10.147 Weak Reaction		< 10	U/mL
Potato Salivary IgA	10.323 Weak Reaction		< 10	U/mL
Potato Salivary IgM	10.147 Weak Reaction		< 10	U/mL
Quinoa Salivary IgA	11.089 Weak Reaction		< 10	U/mL
Quinoa Salivary IgM	10.233 Weak Reaction		< 10	U/mL
Rabbit Salivary IgA	9.630 Negative Reaction		< 10	U/mL
Rabbit Salivary IgM	9.695 Negative Reaction		< 10	U/mL
Rice Salivary IgA	11.200 Weak Reaction		< 10	U/mL
Rice Salivary IgM	9.750 Negative Reaction		< 10	U/mL
Salmon Salivary IgA	10.125 Weak Reaction		< 10	U/mL
Salmon Salivary IgM	9.500 Negative Reaction		< 10	U/mL
Sweet Potato Salivary IgA	10.607 Weak Reaction		< 10	U/mL
Sweet Potato Salivary IgM	9.330 Negative Reaction		< 10	U/mL





HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843
Tel: 714-891-2022; Fax: 714-891-2123
www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Accession No. Test10000	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 08/23/18
-----------------------------------	-------------------------	-----------------------------------	---------------------------	----------------------

Dear Colleague and Owner: Nutriscan Panels #1 and 2 are completed and showed food reactions to: Beef (and Buffalo/Bison), Chicken, Corn (and Cornstarch in supplements), Venison, White Fish (and their Oils), Lentil (including Peas), and Oat. Please avoid all these reactive foods. Best wishes, Jean [W.Jean Dodds, DVM]

Nutriscan test results indicate which of the 24 foods tested are unsuitable for this animal (i.e. tested or above 11.5 units/ml with *either* IgA or IgM antibodies).

Foods testing below 11.5 units/mL should be safe to feed as results were negative or weak (weak reactions reflect background assay "noise").

A **NutriGold** consultation is available, if you have questions or wish advice and suggestions about these results.

Nutriscan Interpretation (see www.nutriscan.org for more details)

Pet should not consume food or treats containing ingredient(s) showing results of **11.5 or greater, that state "Avoid" above, for either IgA and/or IgM antibodies.** Please remember that changing your pet's diet can cause bowel upsets, so gradually changing foods over 7-14 days is preferred. Recommend rechecking salivary food sensitivity or intolerance levels every 6 -12 months. Jean

Degree of reactivity

- **<10 U/mL** indicates a normal food antigen tolerance level =**negative result.**
- **10-11.4 U/mL** indicates a weak reaction; **clinical significance unclear**
- **11.5-11.9 U/mL** indicates a **borderline reaction**
- **12-12.9 U/mL** indicates an **intermediate reaction**
- **13-14.9 U/mL** indicates a **medium reaction**
- **>= 15 U/mL** indicates a **strong reaction**

Differences between antibodies to IgA and IgM in Saliva

Antibodies to IgA measure the immune response to certain foods in secretions like saliva that have occurred over the last 2 years.

Antibodies to IgM measure the body's primary immune response to a recent exposure of certain foods within the last 6 months.



ENRICHING PET LIFE THROUGH DIAGNOSTIC TESTING
world-wide personal mail-in diagnostic lab for vets and pets



HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843

Tel: 714-891-2022; Fax: 714-891-2123

www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Accession No. Test10000	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 08/23/18
-----------------------------------	-------------------------	-----------------------------------	---------------------------	----------------------

Recipes When Some Meats or Fish are Non-Reactive

Suggestions for the diet would be to include only non-reactive meat or fish; vegetables and fruit, with the possible addition of tapioca (cassava root = gluten-free starch). Suggest trying: Chick peas (garbanzo beans), pinto or kidney beans, carrots, zucchini, spinach, green beans, **but no peas or pea fiber (if Lentils are reactive)**, and blueberries, cranberries, pomegranates, (**avoid strawberries**); apples, pears, bananas, and melons (including watermelon). Then, if doing better, try adding tapioca. The dog may also tolerate quail, pheasant, kangaroo, or emu.

If Beef is non-reactive, Buffalo and Bison should be acceptable.

If Lamb is non-reactive, Goat should be OK (including Goat Milk & Yogurt).

Oils – When White-colored Fish and/or Salmon are Reactive

Sources of the important Omega-3 oils are: No menhaden, pollack, sardine or herring oil, if white-colored fish is reactive. No salmon oil, if salmon is reactive.

Krill oil = ancient crustacean (but not with a shellfish allergy).

Plant oils = olive oil, borage oil, sunflower oil, coconut oil (in moderation), primrose oil, moringa oil, but **not canola oil**.

W Jean Daniels, DVM

HEMOLIFE assays and reports use patented non-RIA technology and interpretations. Thyroid 5™ is a trademark.